



PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a preprint version which may differ from the publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/139578>

Please be advised that this information was generated on 2017-12-05 and may be subject to change.

ON THE MATHIEU CONJECTURE FOR $SU(2)$

TEUN DINGS, ERIK KOELINK

ABSTRACT. We study the Mathieu Conjecture for $SU(2)$ using the matrix elements of its unitary irreducible representations. We state a conjecture for the particular case $SU(2)$ implying the Mathieu Conjecture for $SU(2)$.

1. INTRODUCTION

Conjecture 1.1 (Mathieu [6]). *Let G be a compact connected Lie group and let f be complex-valued G -finite function on G such that $\int_G f^P(g) dg = 0$ for every $P \in \mathbb{N}_{>0}$. Then for any complex-valued G -finite function h on G we have $\int_G f^P(g)h(g) dg = 0$ for $P \gg 0$.*

The Mathieu Conjecture 1.1 dates back to 1997 and is closely related to the Jacobian conjecture, since it actually implies the Jacobian conjecture, see [6]. See van den Essen [2], Smale [7] for more information on the history of the Jacobian conjecture. The Mathieu Conjecture 1.1 was proved for abelian compact groups by Duistermaat and Van der Kallen [1] in 1998. We study the Mathieu Conjecture 1.1 for the case $G = SU(2)$. Using explicit formulas for the Haar measure and known representation theoretic properties of $SU(2)$ we make the Mathieu Conjecture 1.1 more explicit. In particular, we use the fact that $SU(2)$ -finite functions are finite linear combinations of matrix elements of finite dimensional irreducible representations of $SU(2)$ and that the matrix elements behave well under a subgroup $K \cong U(1)$ according to suitable characters. Note that the Mathieu Conjecture 1.1 is linear in the G -finite function h , but not in the G -finite function f . By the Peter-Weyl theorem, any $SU(2)$ -finite function is the finite linear combination of matrix elements of irreducible representations. After recalling the necessary results on $SU(2)$ in Section 2, we show in Section 3 the validity of the Mathieu Conjecture 1.1 for f a single matrix element or a sum of two matrix elements. For the sum of three matrix elements there is a partial result. These considerations lead to Conjecture 4.1, and Theorem 4.2 shows that this conjecture implies the Mathieu Conjecture 1.1 for $SU(2)$. Conjecture 4.1 describes the condition $\int_{SU(2)} f(g)^P dg = 0$ for all $P > 0$ in terms of a support condition on the characters of the abelian subgroup $U(1)$ of $SU(2)$ acting from the left and right on the individual matrix elements occurring in f .

We note that the Mathieu Conjecture 1.1 for bi- K -invariant functions is settled by Francoise et al. [3, Cor. 4.1], since the bi- K -invariant $SU(2)$ -finite functions are the polynomials on $[-1, 1]$.

Date: April 17, 2014.

2. $SU(2)$

We briefly recall some required notions of $SU(2)$. Details can be found in e.g. [8], [9]. Let $k(\phi) = \begin{pmatrix} e^{\frac{i}{2}\phi} & 0 \\ 0 & e^{-\frac{i}{2}\phi} \end{pmatrix}$ and $a(\theta) = \begin{pmatrix} \cos \frac{\theta}{2} & i \sin \frac{\theta}{2} \\ i \sin \frac{\theta}{2} & \cos \frac{\theta}{2} \end{pmatrix}$ be elements of $SU(2)$, then any element $g \in SU(2)$ can be expressed in terms of Euler angles $g = k(\phi)a(\theta)k(\psi)$ with $\phi \in [0, 2\pi)$, $\theta \in (0, \pi)$, $\psi \in [-2\pi, 2\pi)$. In terms of the Euler angles the Haar integral is, cf [8, III, §6.1, (5)],

$$\int_{SU(2)} f(g) dg = \frac{1}{16\pi^2} \int_0^{2\pi} \int_0^\pi \int_{-2\pi}^{2\pi} F(\phi, \theta, \psi) \sin \theta d\psi d\theta d\phi, \quad (2.1)$$

where $F(\phi, \theta, \psi) = f(k(\phi)a(\theta)k(\psi))$. Denote the subgroup $K \cong U(1)$ generated by $k(\phi)$. For a function f transforming by a non-trivial K -character under left- or right multiplication by K , we have $\int_{SU(2)} f(g) dg = 0$ by (2.1). The subgroup generated by $a(\theta)$ is the group $SO(2)$.

The finite-dimensional irreducible representations are labeled by the spin $\ell \in \frac{1}{2}\mathbb{N}$ and are of dimension $2\ell + 1$. The standard basis for the representation space is labeled as $\{-\ell, -\ell + 1, \dots, \ell\}$, and the corresponding matrix elements $t_{m,n}^\ell$ are $SU(2)$ -finite functions, and any $SU(2)$ -finite function is a finite linear combination of matrix elements of irreducible finite-dimensional representations. The matrix-elements $t_{m,n}^\ell$ behave well according to left and right action by K , cf. [8, III, §3.3, (3)]

$$t_{m,n}^\ell(k(\phi)g) = e^{-im\phi} t_{m,n}^\ell(g), \quad t_{m,n}^\ell(gk(\psi)) = e^{-in\psi} t_{m,n}^\ell(g). \quad (2.2)$$

In particular, $t_{0,0}^0(g) = 1$, and the algebra of bi- K -invariant $SU(2)$ -finite functions consists of finite linear combinations of $t_{0,0}^\ell$, $\ell \in \mathbb{N}$. For $\ell \in \mathbb{N}$ we have $t_{0,0}^\ell(a(\theta)) = P_\ell(\cos \theta)$, cf. [8, III, §3.9, (5)] where P_ℓ is the Legendre polynomial in its standard normalisation $P_\ell(1) = 1$, [4, §4.5], [5, §1.8.3], which is real-valued on $[-1, 1]$. The Legendre polynomials are orthogonal on $[-1, 1]$ with respect to the uniform measure; $\int_{-1}^1 P_n(x)P_m(x) dx = \delta_{n,m}2/(2n+1)$. Moreover, the Schur orthogonality relations are, [8, III, §6.2, (1)]

$$\int_{SU(2)} t_{m,n}^{\ell_1}(g) \overline{t_{p,q}^{\ell_2}(g)} dg = \frac{1}{2\ell_1+1} \delta_{\ell_1, \ell_2} \delta_{m,p} \delta_{n,q}, \quad (2.3)$$

which in case $m = n = p = q = 0$ give the orthogonality for the Legendre polynomials.

3. THE MATHIEU CONJECTURE FOR $SU(2)$ FOR SIMPLE f

We start using some simple observations related to the condition in the Mathieu Conjecture 1.1 for $G = SU(2)$. Firstly, by the Schur orthogonality relations (2.3)

$$\int_{SU(2)} t_{m,n}^\ell(g) dg \neq 0 \iff \ell = 0. \quad (3.1)$$

Secondly, by the left and right K -behaviour of the matrix elements (2.2) and the Haar measure in Euler angles (2.1) we see

$$\int_{SU(2)} (t_{m_1, n_1}^{\ell_1})^{\alpha_1}(g) \cdots (t_{m_k, n_k}^{\ell_k})^{\alpha_k}(g) dg \neq 0 \implies \sum_{i=1}^k \alpha_i m_i = 0 = \sum_{i=1}^k \alpha_i n_i \quad (3.2)$$

for $\alpha_i \in \mathbb{N}$, $\ell_i \in \frac{1}{2}\mathbb{N}$ and $m_i, n_i \in \{-\ell_i, \dots, \ell_i\}$.

Lemma 3.1. $\int_{SU(2)} (t_{m,n}^\ell)^P(g) dg = 0$ for all integer $P > 0$ if and only if $m \neq 0$ or $n \neq 0$.

Proof. The implication \Leftarrow follows from (3.2). To prove the other implication, we observe that for $\ell \in \mathbb{N}$

$$\int_{SU(2)} (t_{0,0}^\ell(g))^2 dg = \frac{1}{2} \int_0^\pi (P_\ell(\cos \theta))^2 \sin \theta d\theta = \frac{1}{2} \int_{-1}^1 (P_\ell(x))^2 dx > 0. \quad \square$$

Now we can verify the Mathieu Conjecture 1.1 in the case f consists of one matrix element.

Proposition 3.2. *The Mathieu Conjecture 1.1 is true for $G = SU(2)$ with f a single matrix element $f = t_{m,n}^\ell$.*

Proof. Since all non-negative powers of f integrate to zero, Lemma 3.1 shows that $m \neq 0$ or $n \neq 0$, so in particular $\ell \neq 0$. Let $h = t_{a,b}^{\ell_0}$. We assume $m \neq 0$, the case $n \neq 0$ being similar. By (3.2) we see that $Pm + a \neq 0$ implies $\int_{SU(2)} (f(g))^P h(g) dg = 0$, which is the case for $P > |a|/|m|$. \square

The same strategy can also be employed to deal with $f = A_1 t_{m_1, n_1}^{\ell_1} + A_2 t_{m_2, n_2}^{\ell_2}$, where $A_i \in \mathbb{C}$, assuming $A_1 \neq 0 \neq A_2$ and $(\ell_1, m_1, n_1) \neq (\ell_2, m_2, n_2)$. Note

$$\int_{SU(2)} (f(g))^P dg = \sum_{\alpha=0}^P \binom{P}{\alpha} A_1^\alpha A_2^{P-\alpha} \int_{SU(2)} (t_{m_1, n_1}^{\ell_1})^\alpha(g) (t_{m_2, n_2}^{\ell_2})^{P-\alpha}(g) dg. \quad (3.3)$$

Lemma 3.3. *Let f be as above with at least one of (m_1, m_2, n_1, n_2) non-zero, then*

$$\exists P > 0 : \int_{SU(2)} (f(g))^P dg \neq 0 \iff \det \begin{pmatrix} m_1 & m_2 \\ n_1 & n_2 \end{pmatrix} = 0 \wedge m_1 m_2 \leq 0 \wedge n_1 n_2 \leq 0.$$

Remark 3.4. Note that the condition in Lemma 3.3 means that $(0, 0)$ is on the line segment from (m_1, n_1) to (m_2, n_2) .

Proof. \Rightarrow : Since at least one term in the right hand side of (3.3) has to be non-zero, (3.2) shows that $m_1 \alpha + m_2 (P - \alpha) = 0 = n_1 \alpha + n_2 (P - \alpha)$, which gives the result.

\Leftarrow : Note that $\dim \text{Ker} \begin{pmatrix} m_1 & m_2 \\ n_1 & n_2 \end{pmatrix} = 1$. Pick a solution $(\alpha, \beta) \in \mathbb{N}^2$ to $m_1 \alpha + m_2 \beta = 0 = n_1 \alpha + n_2 \beta$, and put $M = \alpha + \beta$. Then

$$\int_{SU(2)} (f(g))^M dg = \binom{\alpha + \beta}{\alpha} A_1^\alpha A_2^\beta \int_{SU(2)} (t_{m_1, n_1}^{\ell_1})^\alpha(g) (t_{m_2, n_2}^{\ell_2})^\beta(g) dg, \quad (3.4)$$

using (3.2), since for $\gamma \neq 0$

$$\begin{pmatrix} m_1 & m_2 \\ n_1 & n_2 \end{pmatrix} \begin{pmatrix} \alpha + \gamma \\ \beta - \gamma \end{pmatrix} = \gamma \begin{pmatrix} m_1 & m_2 \\ n_1 & n_2 \end{pmatrix} \begin{pmatrix} 1 \\ -1 \end{pmatrix} \neq \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

since the kernel is one-dimensional. The integrand on the right hand side of (3.4) is a bi- K -invariant function, so that by (2.1) we can restrict to the integral over $g = a(\theta)$, $\theta \in [0, \pi]$. By [8, III, §3,(3),(4)] the integrand in $a(\theta)$ is real-valued. In case the integral is non-zero

we are done. Otherwise, we put $P = 2M$, and then in the same way there is again at most one non-zero integral in the right hand side of (3.3), namely for $(2\alpha, 2\beta)$. The integral can be restricted to $SO(2)$ as before. Since this is the integral of a square, since the function $(t_{m_1, n_1}^{\ell_1})^\alpha (a(\theta)) (t_{m_2, n_2}^{\ell_2})^\beta (a(\theta))$ is real, the integral is non-zero. \square

Proposition 3.5. *The Mathieu Conjecture 1.1 is true for $G = SU(2)$ with f a sum of two matrix element $f = A_1 t_{m_1, n_1}^{\ell_1} + A_2 t_{m_2, n_2}^{\ell_2}$, where $A_1 \neq 0 \neq A_2$ and $(\ell_1, m_1, n_1) \neq (\ell_2, m_2, n_2)$.*

Proof. It suffices to take $h = t_{a,b}^\ell$ and to assume that $\int_{SU(2)} (f(g))^P dg = 0$ for all $P > 0$. We need to show that $\int_{SU(2)} (f(g))^P t_{a,b}^\ell(g) dg$ vanishes for sufficiently large P .

First assume that not all of m_i 's and n_i 's are zero, then by Lemma 3.3 we have $m_1 m_2 > 0$ or $n_1 n_2 > 0$ or $\det \begin{pmatrix} m_1 & m_2 \\ n_1 & n_2 \end{pmatrix} \neq 0$. Consider the last case, then by (3.2), (3.3) we see that $\int_{SU(2)} (f(g))^P t_{a,b}^\ell(g) dg$ can only be non-zero if

$$m_1 \alpha + m_2 \beta = -a, \quad n_1 \alpha + n_2 \beta = -b, \quad \alpha + \beta = P, \quad \alpha, \beta \in \mathbb{N}.$$

The first two equations have a unique solution $(\alpha_0, \beta_0) \in \mathbb{Q}^2$. In case $(\alpha_0, \beta_0) \in \mathbb{N}^2$, we see that for all $P > \alpha_0 + \beta_0$ the integral is zero. In case $m_1 m_2 > 0$, we consider $m_1 \alpha + m_2 \beta + a = 0$. In case $\text{sgn}(m_1) = \text{sgn}(a)$, we have no solution $(\alpha, \beta) \in \mathbb{N}^2$, so that integral is zero using (3.2), (3.3). In case $\text{sgn}(m_1) = -\text{sgn}(a)$, we see that the integral is zero for $P > |a| / \min(|m_1|, |m_2|)$. The case $n_1 n_2 > 0$ is dealt with analogously.

In case $m_1 = m_2 = n_1 = n_2 = 0$, f is a bi- K -invariant function, and

$$\int_{SU(2)} (f(g))^P dg = \frac{1}{2} \int_{-1}^1 (A_1 P_{\ell_1}(x) + A_2 P_{\ell_2}(x))^P dx.$$

By Boyarchenko's result, see [3, Cor. 4.1], there is no polynomial f such that $\int_{-1}^1 (f(x))^P dx = 0$ for all $P > 0$, so the Mathieu Conjecture 1.1 is trivially valid in this case. \square

The fact that at most one term in the binomial expansion leads to a non-zero integral is typical for f a linear combination of two matrix elements. For a combination of three matrix-elements it gets more complicated.

Proposition 3.6. *Let $f = \sum_{i=1}^3 A_i t_{m_i, n_i}^{\ell_i}$ with $A_i \neq 0$ for all i and let (ℓ_i, m_i, n_i) be mutually different. Assume that $M = \begin{pmatrix} 1 & 1 & 1 \\ m_1 & m_2 & m_3 \\ n_1 & n_2 & n_3 \end{pmatrix}$ has $\text{rank}(M) \neq 2$. Then the Mathieu Conjecture 1.1 is valid for f .*

Proof. The analogue of (3.3) is the trinomial expansion

$$\int_{SU(2)} f^P(g) dg = \sum_{\substack{\alpha_1 + \alpha_2 + \alpha_3 = P \\ \alpha_i \in \mathbb{N}}} \binom{P}{\alpha_1, \alpha_2, \alpha_3} \prod_{i=1}^3 A_i^{\alpha_i} \int_{SU(2)} \prod_{i=1}^3 (t_{m_i, n_i}^{\ell_i})^{\alpha_i}(g) dg. \quad (3.5)$$

As before, it suffices to consider the case $h = t_{a,b}^\ell$. We have to consider the cases $\text{rank}(M) = 1$ and $\text{rank}(M) = 3$. In the first case $m_i = m$ and $n_i = n$ for all i , and the integral in (3.5) is

zero if $m \neq 0$ or $n \neq 0$ by (3.2). In case $m \neq 0$ we see that

$$\int_{SU(2)} f^P(g) t_{a,b}^\ell(g) dg = \sum_{\substack{\alpha_1 + \alpha_2 + \alpha_3 = P \\ \alpha_i \in \mathbb{N}}} \binom{P}{\alpha_1, \alpha_2, \alpha_3} \prod_{i=1}^3 A_i^{\alpha_i} \int_{SU(2)} \prod_{i=1}^3 (t_{m_i n_i}^{\ell_i})^{\alpha_i}(g) t_{a,b}^\ell(g) dg \quad (3.6)$$

can only be non-zero if $Pm + a = 0$, so that for $P > |a|/|m|$ the integral is zero. The case $n \neq 0$ is analogous. In case $m = n = 0$, we see that the condition in the Mathieu Conjecture is not valid using [3, Cor. 4.1] as in the proof of Proposition 3.5.

In case $\text{rank}(M) = 3$, M is invertible with M^{-1} having rational entries. In particular, for each $P \in \mathbb{N}$ there is at most one term in the right hand side of (3.5) which can be non-zero, namely for $\vec{\alpha}_P = \begin{pmatrix} \alpha_1 \\ \alpha_2 \\ \alpha_3 \end{pmatrix} = M^{-1} \begin{pmatrix} P \\ 0 \\ 0 \end{pmatrix}$ under the additional condition $\vec{\alpha}_P \in \mathbb{N}^3$. Assuming that this is the case, we see that, analogous to the proof of Proposition 3.5, $\int_{SU(2)} f^{2P}(g) dg \neq 0$.

So we need to consider the case that $\vec{\alpha}_P \notin \mathbb{N}^3$ for all $P > 0$. Then the integral in (3.6) can only be non-zero in case

$$M^{-1} \begin{pmatrix} P \\ -a \\ -b \end{pmatrix} = \frac{1}{\det(M)} \left(P \begin{pmatrix} m_2 n_3 - m_3 n_2 \\ m_3 n_1 - m_1 n_3 \\ m_1 n_2 - m_2 n_1 \end{pmatrix} - a \begin{pmatrix} n_2 - n_3 \\ n_3 - n_1 \\ n_1 - n_2 \end{pmatrix} - b \begin{pmatrix} m_3 - m_2 \\ m_1 - m_3 \\ m_2 - m_1 \end{pmatrix} \right) \in \mathbb{N}^3$$

Since $\vec{\alpha}_P$ corresponds to the first term, i.e. $a = b = 0$, and $\vec{\alpha}_P \notin \mathbb{N}^3$ for all $P > 0$ we have $\det(M)^{-1}(m_i n_{i+1} - m_{i+1} n_i) < 0$ for some $i \in \{1, 2, 3\}$ with convention $m_4 = m_1$, $n_4 = n_1$. Then for $P > (|a||n_i - n_{i+1}| + |b||m_{i+1} - m_i|)/|m_i n_{i+1} - m_{i+1} n_i|$ the i -th coefficient is negative, so that the integral in (3.6) is zero. \square

Remark 3.7. In case $\text{rank}(M) = 1$ the convex hull C of $((m_i, n_i))_{i=1}^3$ equals $\{(m, n)\}$, and in case $\text{rank}(M) = 3$ we have $(0, 0) \in C$ if and only if $\exists \vec{\alpha} \in \mathbb{Q}_{\geq 0}^3$ with $M\vec{\alpha} = (1, 0, 0)^t$. From the proof of Proposition 3.6 we see that $\int_{SU(2)} f(g)^P dg = 0$ for all $P > 0$ precisely when $(0, 0) \notin C$ in the cases $\text{rank}(M) \neq 2$. In case $\text{rank}(M) = 2$ the integral in (3.5) can have more than one non-zero term, and we have no control on possible cancellations. However, one expects that these cancellations cannot occur for all multiples of P as well. The techniques of Francoise et al. [3] might be useful in this regard considering it as polynomial identities in the A_i 's.

4. AN ALTERNATIVE CONJECTURE FOR THE MATHIEU CONJECTURE FOR $SU(2)$

Consider an arbitrary $SU(2)$ -finite function $f = \sum_{i=1}^k A_i t_{m_i, n_i}^{\ell_i}$ with $A_i \neq 0$ for every $1 \leq i \leq k$, then applying the multinomial theorem shows that if

$$\int_{SU(2)} (f(g))^P dg = \sum_{\alpha_i \in \mathbb{N}, \sum_{i=1}^k \alpha_i = P} \binom{P}{\alpha_1, \dots, \alpha_k} \prod_{i=1}^k A_i^{\alpha_i} \int_{SU(2)} \prod_{i=1}^k (t_{m_i, n_i}^{\ell_i})^{\alpha_i}(g) dg \neq 0 \quad (4.1)$$

for some $P > 0$, then for some $(\alpha_1, \dots, \alpha_k)$ we have $\sum_{i=1}^k \frac{\alpha_i}{P} m_i = \sum_{i=1}^k \frac{\alpha_i}{P} n_i = 0$ by (3.2), so $(0, 0)$ is in the convex hull C of $((m_i, n_i))_{i=1}^k$ over \mathbb{Q} .

Conjecture 4.1. *For any $SU(2)$ -finite function $f = \sum_{i=1}^k A_i t_{m_i, n_i}^{\ell_i}$, $A_i \neq 0$ for all $1 \leq i \leq k$, we have that $\int_{SU(2)} (f(g))^P dg = 0$ for all $P \in \mathbb{N}_{>0}$ if and only if $(0, 0)$ is not contained in the closed convex hull of $((m_i, n_i))_{i=1}^k$.*

Lemma 3.1, Remarks 3.4, 3.7 support Conjecture 4.1.

Theorem 4.2. *Assume Conjecture 4.1 holds, then the Mathieu Conjecture 1.1 for $SU(2)$ holds.*

Proof. It suffices to show that $\int_{SU(2)} (f(g))^P t_{a,b}^{\ell}(g) dg = 0$ for P sufficiently large assuming that $(0, 0)$ is not contained in the closed convex hull C of $((m_i, n_i))_{i=1}^k$. Using (4.1) we see that $\int_{SU(2)} (f(g))^P t_{a,b}^{\ell}(g) dg$ can only be non-zero if $(-\frac{a}{P}, -\frac{b}{P}) \in C$. Since $(0, 0) \notin C$, we see that for P sufficiently large this is not the case and the integral is zero. \square

Acknowledgement. TD thanks Wilberd van der Kallen, Hjalmar Rosengren, Harm Derksen, Pablo Román for useful discussions. We thank Harm Derksen for pointing out Boyarchenko's result and Arno van den Essen for pointing out [3].

REFERENCES

- [1] J.J. Duistermaat, W. van der Kallen, Constant terms in powers of a Laurent polynomial, *Indag. Math. (N.S.)* **9** (1998), 221–231.
- [2] A. van den Essen, *Polynomial automorphisms and the Jacobian conjecture*, Progress in Math. 190, Birkhäuser, 2000.
- [3] J.P. Francoise, F. Pakovich, Y. Yomdin, W. Zhao, Moment vanishing problem and positivity: some examples, *Bull. Sci. math.* **135** (2011), 10–32.
- [4] M.E.H. Ismail, *Classical and quantum orthogonal polynomials in one variable*, Cambridge Univ. Press, 2009.
- [5] R. Koekoek, R.F. Swarttouw, The Askey-scheme of hypergeometric orthogonal polynomials and its q -analogue, online at <http://aw.twi.tudelft.nl/~koekoek/askey.html>, Report 98-17, Technical University Delft, 1998.
- [6] O. Mathieu, Some conjectures about invariant theory and their applications, pp. 263–279 in *Algèbre non commutative, groupes quantiques et invariants* (eds. J. Alev, G. Cauchon), Séminaires et Congrès **2**, Soc. Math. France, 1997.
- [7] S. Smale, Mathematical problems for the next century. *Math. Intelligencer* **20** (1998), 7–15.
- [8] N.Ja. Vilenkin, *Special functions and the theory of group representations*, Translations Math. Monographs 22, AMS, 1968.
- [9] N.Ja. Vilenkin, A.U. Klimyk, *Representation of Lie groups and special functions* Vol. 1. Simplest Lie groups, special functions and integral transforms, Math. and its Appl. 72, Kluwer, 1991.

RADBOUD UNIVERSITEIT, IMAPP, HEYENDAALSEWEG 135, 6525 GL NIJMEGEN, THE NETHERLANDS
E-mail address: teundings@gmail.com, e.koelink@math.ru.nl